

ZETA FUNCTION ON SURFACES OF REVOLUTION

THALIA D. JEFFRES, KLAUS KIRSTEN, AND TIANSHI LU

ABSTRACT. In this paper we applied the contour integral method for the zeta function associated with a differential operator to the Laplacian on a surface of revolution. Using the WKB expansion, we calculated the residues and values of the zeta function at several important points. The results agree with those obtained from the heat kernel expansion. We also obtained a closed form formula for the determinant of the Laplacian on such a surface.

1. INTRODUCTION

Spectral functions, namely functions of the spectrum of, usually, second order differential operators play an important role in mathematics and physics. The possibly most important such spectral function is the zeta function which directly relates to topics such as analytic torsion [40], the heat kernel [27, 43], Casimir energy [4, 5, 13, 20, 36], effective actions and quantum tunneling [7, 11, 14, 15, 16, 21, 28, 42] and the study of critical metrics [10, 38, 39]. The relationship between the zeta function and all these topics is established by the fact that very specific properties of the zeta function encode the information needed in these areas. For example residues and values at specific points contain information about the short time heat kernel asymptotics with all the geometric information it contains, the (finite part of its) value at $s = -1/2$ relates to the Casimir energy, and its derivative at $s = 0$ is relevant in all the other subjects mentioned above; see, e.g., [29]. It is therefore no surprise that an enormous effort is made to understand spectral zeta functions as completely as possible.

The situation is particularly satisfying for the one-dimensional case where functional determinants can be obtained in closed form as boundary values of a suitable initial value problem [8, 9, 17, 22, 23, 26, 34]. A contour integral approach established in [31, 32] was used to rederive this case and generalize it to arbitrary boundary conditions. This approach has the advantage that it allows for the systematic study of higher dimensional configurations as long as the Laplacian separates in a suitable fashion. Examples are Laplacians with spherically symmetric potentials [18, 19] and Laplacians for spherical suspensions [24]. In those cases the analysis of the associated zeta function is based on explicitly known special functions, namely Bessel functions and Legendre functions.

Another example, studied in the present paper, is the Laplacian on a compact surface of revolution. These surfaces are geometrically interesting enough to exhibit many features that appear in the study of spectral functions and their asymptotics, while at the same time the rotational symmetry allows the explicit calculations that are necessary in order to understand meromorphic extensions of the zeta function to regions outside of the domain of convergence. The complication with this example is that no known special function emerges as a solution to the relevant initial value problem because the function f whose graph is used to generate the surface of revolution is kept fairly general. As a result needed asymptotic behavior has to be determined from scratch starting from an ordinary differential equation. The WKB method is our method of choice [3, 35] and it allows us to obtain the values or residues of the

zeta function at several important points, in addition to the determinant of the Laplacian. The relation between the values of the zeta function and the global geometrical properties of the surface is revealed. Furthermore it is notable that the result for the determinant is a closed-form formula.

The structure of the paper is as follows. In Section 2 we present the geometric preliminaries, define the generalized zeta function, and reformulate it as sum of contour integrals. The zeta function is written as the sum of a one-dimensional term and a two-dimensional term, which are calculated in Section 3 and Section 4 respectively. The values and residues of the zeta function at several selected points are calculated along with the determinant of the Laplacian in Section 4. We conclude in Section 5 with a comparison of our method and other approaches.

2. ZETA FUNCTION FOR A SURFACE OF REVOLUTION

2.1. Laplacian on a surface of revolution. We consider the Laplacian on a surface of revolution with Dirichlet boundary values. Let $f \in C^2(a, b)$ be a positive function from $[a, b]$ to \mathbb{R} , and let M be the compact surface with boundary that is generated by revolving the graph of f around the x -axis. With the metric induced by the Euclidean metric of \mathbb{R}^3 , (M, g) becomes a Riemannian manifold. In the natural coordinates (x, θ) , the metric is

$$\begin{bmatrix} 1 + f'(x)^2 & 0 \\ 0 & f^2(x) \end{bmatrix}.$$

Recalling the formula for the Christoffel symbols in terms of the metric,

$$\Gamma_{ij}^k = \frac{1}{2} \sum_{l=1}^2 g^{kl} \left(\frac{\partial g_{il}}{\partial x_j} + \frac{\partial g_{jl}}{\partial x_i} - \frac{\partial g_{ij}}{\partial x_l} \right),$$

we have

$$\Gamma_{11}^1 = \frac{f'f''}{1+f'^2}, \quad \Gamma_{12}^2 = \Gamma_{21}^2 = \frac{f'}{f}, \quad \Gamma_{22}^1 = \frac{-ff'}{1+f'^2},$$

and the others are zero. We then calculate that

$$\Delta u = \frac{1}{1+f'^2} \left[\frac{\partial^2 u}{\partial x^2} + \left(\frac{f'}{f} - \frac{f'f''}{1+f'^2} \right) \frac{\partial u}{\partial x} + \frac{1+f'^2}{f^2} \frac{\partial^2 u}{\partial \theta^2} \right]. \quad (1)$$

The eigenvalue equation is $\Delta u = -\lambda u$. The following facts are standard [25, 41]:

- (1) The eigenvalues are real and positive; if labeled and ordered as $0 < \lambda_1 \leq \lambda_2 \leq \dots$, then $\lambda_k \rightarrow \infty$.
- (2) For each λ_k , the corresponding eigenspace in $L^2(M)$ is finite-dimensional.
- (3) The spectrum of M is discrete.

We remark here that zero is not an eigenvalue of the Laplacian if Dirichlet boundary values are taken.

2.2. Generalized zeta function. Our spectral function for the Laplacian generalizes the well-known Riemann zeta function,

$$\zeta_R(s) = \sum_{k=1}^{\infty} k^{-s}, \quad (2)$$

which converges for $\Re(s) > 1$. For the Laplacian on a surface of revolution with Dirichlet boundary condition, if the eigenvalues are labeled and ordered as $0 < \lambda_1 \leq \lambda_2 \leq \dots$, the generalized zeta function is defined to be

$$\zeta(s) = \sum_{\lambda_k \in \text{Spec}(\Delta)} \lambda_k^{-s}, \quad (3)$$

for those values of $s \in \mathbb{C}$ for which this sum converges. Weyl's asymptotic formula can be used to determine the domain of convergence of Eq. (3) [12]. On a compact Riemannian manifold M of dimension n ,

$$\lambda_k^{\frac{n}{2}} \sim \frac{2^n \pi^{\frac{n}{2}} \Gamma(\frac{n}{2} + 1)}{\text{Vol}(M)} k, \quad k \rightarrow \infty.$$

From this we determine that the zeta function is defined for complex values of s having $\Re(s) > n/2$, in our case $\Re(s) > 1$. However, just as for the Riemann zeta function, much information of geometric and physical significance is contained in $\zeta(s)$ for complex values of s outside of this region. $\zeta(s)$ has a simple pole at $s = 1$. The holomorphic continuation of $\zeta(s)$ to $\Re(s) < 1$ turns out to have simple poles at $s = \frac{1}{2} - n$, $n \in \mathbb{N}$. Here we obtain expressions for the residues of $\zeta(s)$ at $s = 1, \frac{1}{2}, -\frac{1}{2}$, and its value at $s = 0$, in terms of geometrical properties of the surface. Furthermore we compute the determinant of the Laplacian using the relation with the zeta function given by [16, 28, 40]

$$\ln \det \Delta = -\zeta'(0). \quad (4)$$

We obtain a formula for $\zeta'(0)$ given entirely in terms of the function f whose graph generates the surface of revolution.

2.3. Reformulation of the zeta function as a contour integral. A fundamental difficulty in studying generalized zeta functions is that the eigenvalues are not known explicitly except for special cases. However, as described in Ref. [29, 30], the zeta function can be reformulated as a contour integral, which is highly suitable to such investigations. Here we adapt this method to the Laplacian on a surface of revolution.

The rotational symmetry of M suggests the use of separation of variables to describe the eigenvalues of the Laplacian. The eigenfunctions of the Laplacian on the cross-section S^1 , denoted by $u_k(\theta)$, $k \in \mathbb{Z}$, are $e^{ik\theta}$, with eigenvalue $-k^2$. Let $u(x, \theta) = \phi(x)u_k(\theta)$. By Eq. (1), u is an eigenfunction of Δ on M with eigenvalue λ if and only if ϕ is a nontrivial solution to

$$\phi''(x) + \left(\frac{f'}{f} - \frac{f'f''}{1+f'^2} \right) \phi'(x) + \left(\lambda - \frac{k^2}{f^2} \right) (1 + f'^2) \phi(x) = 0, \quad (5)$$

satisfying the boundary conditions

$$\phi(a) = 0, \quad \phi(b) = 0.$$

Denote by $\phi_k(\lambda; x)$ the solution to the initial value problem consisting of the same differential equation (5), but with initial conditions

$$\phi(a) = 0, \quad \phi'(a) = 1. \quad (6)$$

Then for each integer k , $\lambda \in \mathbb{C}$ is an eigenvalue if and only if $\phi_k(\lambda; b) = 0$. For each fixed integer k , order the eigenvalues and denote them by $0 < \lambda_{k,1} \leq \lambda_{k,2} \leq \dots$. Since $\phi_k(\lambda; b)$,

regarded as a function of λ , vanishes at and only at $\lambda_{k,n}$, $n \in \mathbb{Z}^+$, its canonical product representation [1] is

$$\phi_k(\lambda; b) = \phi_k(0; b) \prod_{n=1}^{\infty} \left(1 - \frac{\lambda}{\lambda_{k,n}}\right).$$

The convergence of the infinite product is guaranteed by Weyl's asymptotic formula. Let

$$D_k(\lambda) = \frac{\phi_k(\lambda; b)}{\phi_k(0; b)} = \prod_{n=1}^{\infty} \left(1 - \frac{\lambda}{\lambda_{k,n}}\right).$$

By the Residue Theorem, we may express $\zeta(s)$ as the sum of contour integrals.

$$\zeta(s) = \frac{1}{2\pi i} \int_{\Gamma} \lambda^{-s} \frac{d \ln D_0(\lambda)}{d\lambda} d\lambda + 2 \sum_{k=1}^{\infty} \frac{1}{2\pi i} \int_{\Gamma} \lambda^{-s} \frac{d \ln D_k(\lambda)}{d\lambda} d\lambda, \quad (7)$$

where Γ is any counterclockwise contour that encloses all the $\lambda_{k,n}$'s but not the origin. The factor 2 in the second term above comes from the fact that $\lambda_{k,n} = \lambda_{-k,n}$. Following Ref. [30] (Fig. 3), the contour can be chosen to go around the negative axis, from $-\infty$ to 0 above the real axis, along which $\lambda = e^{i\pi}y$, and from 0 to $-\infty$ below the real axis, along which $\lambda = e^{-i\pi}y$. Using this contour, Eq. (7) is simplified to be

$$\zeta(s) = \zeta_1(s) + \zeta_2(s), \quad (8)$$

where

$$\zeta_1(s) = \frac{\sin \pi s}{\pi} \int_0^{\infty} y^{-s} \frac{d \ln D_0(-y)}{dy} dy, \quad (9)$$

$$\zeta_2(s) = 2 \frac{\sin \pi s}{\pi} \sum_{k=1}^{\infty} \int_0^{\infty} y^{-s} \frac{d \ln D_k(-y)}{dy} dy. \quad (10)$$

Note, that Eq. (10) should be understood in the sense that the analytical continuation of the integral is constructed first and the summation of that result is performed afterwards.

Next, in Section 3, we will use the WKB method to solve Eq. (5) with $k = 0$, and compute important values of $\zeta_1(s)$ and $\zeta'_1(0)$. In Section 4 we will adapt the method to the series over k , and compute $\zeta_2(s)$ along with $\zeta'_2(0)$.

3. ZETA FUNCTION AND DERIVATIVE WITH $k = 0$

By Weyl's asymptotic formula Eq. (9) converges for $\Re s > \frac{1}{2}$. To obtain the meromorphic extension of $\zeta_1(s)$ to $\Re s < \frac{1}{2}$, we apply the WKB method to determine the asymptotics of $\phi_0(-y; b)$; see for example [3, 35] for a discussion of the WKB method. First, we make the change of variables $y = z^2$ in the differential equation (5) with $k = 0$. It becomes

$$\phi''(x) + \left(\frac{f'}{f} - \frac{f'f''}{1+f'^2} \right) \phi'(x) - z^2(1+f'^2)\phi(x) = 0. \quad (11)$$

Denote the coefficient of $\phi'(x)$ in Eq. (11) by $u(x)$,

$$u(x) = \frac{f'}{f} - \frac{f'f''}{1+f'^2}.$$

Let

$$\phi(x) = \exp \left(\int_a^x [S(z; \sigma) - \frac{u(\sigma)}{2}] d\sigma \right). \quad (12)$$

Substituted into Eq. (11), the equation for $S(z; x)$ is

$$S'(z; x) + S(z; x)^2 = (1 + f'^2)z^2 + \frac{u^2}{4} + \frac{u'}{2},$$

where the prime indicates differentiation with respect to x . The solution can be written as asymptotic expansions in z . It has two branches, namely,

$$S_{\pm}(z; x) = \pm s_{-1}z + s_0 \pm \frac{s_1}{z} + \dots = \pm(s_{-1}z + \frac{s_1}{z} + \dots) + (s_0 + \frac{s_2}{z^2} + \dots), \quad (13)$$

where the coefficients $s_n(x)$ can be computed recursively. The first three coefficients are

$$s_{-1} = \sqrt{1 + f'^2}, \quad s_0 = -\frac{f'f''}{2(1 + f'^2)}, \quad s_1 = -\frac{1}{8} \frac{f'^2}{f^2 \sqrt{1 + f'^2}} + \frac{1}{4} \frac{f''}{f(1 + f'^2)^{3/2}}. \quad (14)$$

Applying the initial condition $\phi(a) = 0$ and $\phi'(a) = 1$, the solution to Eq. (11) is

$$\begin{aligned} \phi_0(-z^2; x) &= e^{-\frac{1}{2} \int_a^x u(\sigma) d\sigma} \frac{e^{\int_a^x S_+(z; \sigma) d\sigma} - e^{\int_a^x S_-(z; \sigma) d\sigma}}{S_+(z; a) - S_-(z; a)} \\ &= e^{\int_a^x [S_+(z; \sigma) - \frac{1}{2} u(\sigma)] d\sigma} \frac{1 - e^{-\int_a^x [S_+(z; \sigma) - S_-(z; \sigma)] d\sigma}}{S_+(z; a) - S_-(z; a)}. \end{aligned}$$

Let $S_{\pm} = \pm S_1 + S_2$, where

$$S_1(z; x) = s_{-1}z + \frac{s_1}{z} + \dots, \quad S_2(z; x) = s_0 + \frac{s_2}{z^2} + \dots. \quad (15)$$

Notice that

$$S'_+ + S_+^2 = S'_- + S_-^2 \Rightarrow S'_1 + 2S_1S_2 = 0 \Rightarrow \ln S_1(z; a) - \ln S_1(z; b) = 2 \int_a^b S_2(z; x) dx.$$

$\phi_0(-z^2; b)$ can be written as

$$\phi_0(-z^2; b) = e^{\int_a^b [S_1(z; x) - \frac{1}{2} u(x)] dx} \frac{1 - e^{-\int_a^b 2S_1(z; x) dx}}{2\sqrt{S_1(z; a)S_1(z; b)}}. \quad (16)$$

Now we apply Eq. (16) to the contour integral Eq. (9). Noticing that Eq. (13) is invalid as $z \rightarrow 0$, we write the zeta function as

$$\zeta_1(s) = \frac{\sin \pi s}{\pi} \left[\int_0^1 y^{-s} \frac{d \ln D_0(-y)}{dy} dy + \int_1^\infty z^{-2s} \frac{d \ln D_0(-z^2)}{dz} dz \right]. \quad (17)$$

By Eq. (16),

$$\begin{aligned} \ln D_0(-z^2) &= \int_a^b \left[S_1(z; x) - \frac{u(x)}{2} \right] dx - \frac{\ln S_1(z; a) + \ln S_1(z; b)}{2} \\ &\quad + \ln \left[1 - e^{-2 \int_a^b S_1(z; x) dx} \right] - \ln(2\phi_0(0; b)). \end{aligned} \quad (18)$$

To compute $dD_0(-z^2)/dz$, we need the asymptotic expansion of $\partial S_1(z; x)/\partial z$, which can be derived from Eq. (15), and the expansion of $\partial \ln S_1(z; x)/\partial z$. Let

$$\frac{\partial \ln S_1(z; x)}{\partial z} = \frac{\partial}{\partial z} \left[\ln(s_{-1}z) + \frac{s_1}{s_{-1}z^2} + \dots \right] = \frac{t_0(x)}{z} + \frac{t_1(x)}{z^3} + \dots,$$

where

$$t_0(x) = 1, \quad t_1(x) = -2 \frac{s_1}{s_{-1}} = \frac{1}{4} \frac{f'^2}{f^2(1+f'^2)} - \frac{1}{2} \frac{f''}{f(1+f'^2)^2}. \quad (19)$$

Substituted into Eq. (18),

$$\begin{aligned} \frac{\partial}{\partial z} \ln D_0(-z^2) &= \int_a^b s_{-1}(x) dx - \frac{1}{z^2} \int_a^b s_1(x) dx - \dots \\ &\quad - \frac{1}{2} \frac{t_0(a) + t_0(b)}{z} - \frac{1}{2} \frac{t_1(a) + t_1(b)}{z^3} - \dots \end{aligned}$$

Now we substitute the equation above into the zeta function in Eq. (17). The last integral above is uniformly bounded for s with $\Re s$ bounded below, while the integration from 0 to 1 in Eq. (17) is uniformly bounded for s bounded above by $1 - \epsilon$ for any $\epsilon > 0$. Using the meromorphic extension of the following integrals to $s \in \mathbb{C}$,

$$\int_1^\infty z^{-2s} z^{-n} dz = \frac{1}{2s + n - 1}, \quad n = 0, 1, 2, \dots \quad (20)$$

we can compute the residue of the integral from 1 to ∞ in Eq. (17) at $s = \frac{1}{2}, 0, -\frac{1}{2}$, etc. If s is a half integer, $\sin \pi s \neq 0$, and ζ_1 has a single pole at s for a general $f(x)$. If s is an integer, $\sin \pi s$ has a single zero at s , and so ζ_1 has a finite value at s . For example, using Eq. (20) with $n = 0, 1, 2, 3$ respectively, and plugging into Eqs. (14) and (19), we have

$$\text{Res}_{\zeta_1} \left(\frac{1}{2} \right) = \frac{1}{2\pi} \int_a^b s_{-1}(x) dx = \frac{1}{2\pi} \int_a^b \sqrt{1 + f'(x)^2} dx, \quad (21)$$

$$\zeta_1(0) = -\frac{t_0(a) + t_0(b)}{4} = -\frac{1}{2}, \quad (22)$$

$$\text{Res}_{\zeta_1} \left(-\frac{1}{2} \right) = \frac{1}{2\pi} \int_a^b s_1(x) dx = \frac{1}{2\pi} \int_a^b \left[-\frac{1}{8} \frac{f'^2}{f^2 \sqrt{1 + f'^2}} + \frac{1}{4} \frac{f''}{f(1 + f'^2)^{3/2}} \right] dx, \quad (23)$$

$$\begin{aligned} \zeta_1(-1) &= \frac{t_1(a) + t_1(b)}{4} = \frac{1}{16} \left[\frac{f'(a)^2}{f(a)^2(1 + f'(a)^2)} - \frac{2f''(a)}{f(a)(1 + f'(a)^2)^2} \right. \\ &\quad \left. + \frac{f'(b)^2}{f(b)^2(1 + f'(b)^2)} - \frac{2f''(b)}{f(b)(1 + f'(b)^2)^2} \right]. \end{aligned} \quad (24)$$

The calculation can be continued for $s < -1$, but the expressions become very complicated and we do not include them here.

Derivative of $\zeta_1(s)$ at $s = 0$

To compute $\zeta_1'(0)$, we use integration by parts. Rewrite $\zeta_1(s)$ as

$$\begin{aligned} \zeta_1(s) &= \frac{\sin \pi s}{\pi} s \int_0^\infty y^{-s-1} \ln D_0(-y) dy \\ &= \frac{\sin \pi s}{\pi} s \left[\int_0^1 y^{-s-1} \ln D_0(-y) dy + 2 \int_1^\infty z^{-2s-1} \ln D_0(-z^2) dz \right]. \end{aligned} \quad (25)$$

Since $\ln D_0(-y) = O(y)$ for $y \rightarrow 0$, the integration from 0 to 1 is bounded for $s = 0$. For the integration from 1 to ∞ , by the asymptotic expansion Eq. (15),

$$S_1(z; x) = s_{-1}(x)z + O(z^{-1}) = s_{-1}(x)z \cdot [1 + O(z^{-2})],$$

as $z \rightarrow \infty$. Substituted into Eq. (18),

$$\begin{aligned} \ln D_0(-z^2) &= \int_a^b \left[s_{-1}(x)z + O(z^{-1}) - \frac{u(x)}{2} \right] dx - \frac{1}{2} \ln(s_{-1}(a)z) - \frac{1}{2} \ln(s_{-1}(b)z) \\ &\quad - O(z^{-2}) + \ln \left[1 - \exp(-2 \int_a^b (s_{-1}(x)z + O(z^{-1})) dx) \right] - \ln(2\phi_0(0; b)) \\ &= \int_a^b \left[s_{-1}(x)z - \frac{u(x)}{2} \right] dx - \ln z - \frac{\ln s_{-1}(a) + \ln s_{-1}(b)}{2} - \ln(2\phi_0(0; b)) + O(z^{-1}). \end{aligned}$$

Substituted into Eq. (25), as $s \rightarrow 0$,

$$\begin{aligned} \zeta_1(s) &= 2s \frac{\sin \pi s}{\pi} \int_1^\infty z^{-2s-1} \ln D_0(-z^2) dz \\ &= 2s^2 \int_1^\infty z^{-2s-1} \left\{ z \int_a^b s_{-1}(x) dx - \ln z \right\} dz + O(s^2) \\ &\quad - 2s^2 \int_1^\infty z^{-2s-1} dz \left\{ \frac{1}{2} \int_a^b u(x) dx + \frac{\ln s_{-1}(a) + \ln s_{-1}(b)}{2} + \ln(2\phi_0(0; b)) \right\} \\ &= \frac{2s^2}{2s-1} \int_a^b \sqrt{1+f'^2} dx - \frac{1}{2} + O(s^2) \\ &\quad - s \left\{ \frac{1}{2} \int_a^b u(x) dx + \frac{\ln s_{-1}(a) + \ln s_{-1}(b)}{2} + \ln(2\phi_0(0; b)) \right\}. \end{aligned}$$

This gives $\zeta_1(0) = -\frac{1}{2}$, as in Eq. (22), and furthermore the following theorem.

Theorem 3.1. *The derivative of the zeta function in the one-dimensional case is given by*

$$\zeta_1'(0) = -\frac{\ln f(a) + \ln f(b)}{2} - \ln 2A, \quad (26)$$

where

$$A = \int_a^b \frac{\sqrt{1+f'(x)^2}}{f(x)} dx. \quad (27)$$

Proof. By the calculations above, we have

$$\zeta_1'(0) = - \left(\frac{1}{2} \int_a^b u(x) dx + \frac{\ln s_{-1}(a) + \ln s_{-1}(b)}{2} + \ln(2\phi_0(0; b)) \right).$$

$\phi_0(0; b)$ can be computed exactly,

$$\phi_0(0; b) = \frac{f(a)}{\sqrt{1+f'(a)^2}} A,$$

where A is defined by Eq. (27). Combined with $u(x) = (\ln f - \ln \sqrt{1+f'^2})'$, Eq. (26) follows from a simplification. \square

4. FULL ZETA FUNCTION AND DERIVATIVE

For $k \neq 0$, we make the change of variables $y = k^2 z$. The differential equation for $\phi(x)$ becomes

$$\phi''(x) + u(x)\phi'(x) - k^2(1 + f'^2) \left(z + \frac{1}{f^2} \right) \phi(x) = 0, \quad (28)$$

where $u(x)$ is as defined in Section 3. The zeta function in Eq. (10) becomes

$$\zeta_2(s) = 2 \frac{\sin \pi s}{\pi} \sum_{k=1}^{\infty} k^{-2s} \int_0^{\infty} z^{-s} \frac{d \ln D_k(-k^2 z)}{dz} dz. \quad (29)$$

Because the integrand depends on k , the sum and the integral do not factor, but the sum nevertheless resembles the Riemann zeta function. This is a principal motivation for the change of variable. We define $S(k, z; x)$ as in Eq. (12), and Eq. (16) holds with $\phi_0(-z^2; b)$ replaced by $\phi_k(-k^2 z; b)$. The equation for $S(k, z; x)$ is

$$S'(k, z; x) + S(k, z; x)^2 = \frac{1 + f'^2}{f^2} (zf^2 + 1)k^2 + \frac{u^2}{4} + \frac{u'}{2},$$

and the solutions are written as asymptotic expansions in k ,

$$S_1(k, z; x) = s_{-1}(z; x)k + \frac{s_1(z; x)}{k} + \dots \quad (30)$$

The first two coefficients are

$$s_{-1}(z; x) = \frac{\sqrt{1 + f'^2}}{f} \sqrt{t + 1}, \quad s_1(z; x) = \frac{t}{(t + 1)^{3/2}} \left[-\frac{1}{8} \frac{t - 4}{t + 1} \frac{f'^2}{f \sqrt{1 + f'^2}} + \frac{1}{4} \frac{f''}{(1 + f'^2)^{3/2}} \right], \quad (31)$$

where

$$t(x, z) = zf(x)^2. \quad (32)$$

Recall that $D_k(-k^2 z) = \phi_k(-k^2 z; b)/\phi_k(0; b)$. By Eq. (18),

$$\begin{aligned} \ln D_k(-k^2 z) &= \int_a^b \left[S_1(k, z; x) - \frac{u(x)}{2} \right] dx - \frac{\ln S_1(k, z; a) + \ln S_1(k, z; b)}{2} \\ &\quad + \ln \left[1 - e^{-2 \int_a^b S_1(k, z; x) dx} \right] - \ln(2\phi_k(0; b)). \end{aligned} \quad (33)$$

Similarly, let

$$\frac{\partial \ln S_1(k, -k^2 z; x)}{\partial z} = f^2 \frac{\partial}{\partial t} \left[\ln(s_{-1}k) + \frac{s_1}{s_{-1}} k^{-2} + \dots \right] = f^2 \left[t_0(z; x) + \frac{t_1(z; x)}{k^2} + \dots \right],$$

where

$$t_0 = \frac{1}{2(1 + t)}, \quad t_1 = \frac{t^2 - 10t + 4}{8(1 + t)^4} \frac{f'^2}{1 + f'^2} + \frac{1 - t}{4(1 + t)^3} \frac{ff''}{(1 + f'^2)^2}. \quad (34)$$

Substituted into Eq. (33),

$$\begin{aligned} \frac{\partial}{\partial z} \ln D_k(-k^2 z) &= k \int_a^b f(x)^2 \frac{\partial s_{-1}(z; x)}{\partial t} dx + k^{-1} \int_a^b f(x)^2 \frac{\partial s_1(z; x)}{\partial t} dx + \dots \\ &\quad - \frac{t_0(z; a)f(a)^2 + t_0(z; b)f(b)^2}{2} - \frac{t_1(z; a)f(a)^2 + t_1(z; b)f(b)^2}{2k^2} - \dots \\ &\quad + \frac{2e^{-2 \int_a^b S_1(k, z; x) dx}}{1 - e^{-2 \int_a^b S_1(k, z; x) dx}} \frac{\partial}{\partial z} \int_a^b S_1(k, z; x) dx. \end{aligned}$$

Substituted into Eq. (29),

$$\begin{aligned} \zeta_2(s) &= 2 \frac{\sin \pi s}{\pi} \zeta_R(2s-1) \int_a^b dx f(x)^{2s} \int_0^\infty t^{-s} \frac{\partial}{\partial t} s_{-1}(z; x) dt \\ &\quad + 2 \frac{\sin \pi s}{\pi} \zeta_R(2s+1) \int_a^b dx f(x)^{2s} \int_0^\infty t^{-s} \frac{\partial}{\partial t} s_1(z; x) dt + \dots \\ &\quad - \frac{\sin \pi s}{\pi} \zeta_R(2s) \int_0^\infty [t_0(z; a)f(a)^{2s} t^{-s} + t_0(z; b)f(b)^{2s} t^{-s}] dt \\ &\quad - \frac{\sin \pi s}{\pi} \zeta_R(2s+2) \int_0^\infty [t_1(z; a)f(a)^{2s} t^{-s} + t_1(z; b)f(b)^{2s} t^{-s}] dt - \dots \\ &\quad + 2 \frac{\sin \pi s}{\pi} \sum_{k=1}^\infty k^{-2s} \int_0^\infty dz z^{-s} \frac{2e^{-2 \int_a^b S_1(k, z; x) dx}}{1 - e^{-2 \int_a^b S_1(k, z; x) dx}} \frac{\partial}{\partial z} \int_a^b S_1(k, z; x) dx. \end{aligned}$$

As before, ζ_R is the Riemann zeta function. Since $S_1(k, z; x) = O(k\sqrt{z})$ for large k or z , the last term above does not contribute to the residues or values of $\zeta_2(s)$ for s being integers or half integers below (including) 1. Using the following standard formulas [33],

$$\frac{\sin \pi s}{\pi} = \frac{1}{\Gamma(s)\Gamma(1-s)},$$

and

$$\int_0^\infty \frac{z^{-s}}{(z+1)^p} dz = B(1-s, s+p-1) = \frac{\Gamma(1-s)\Gamma(s+p-1)}{\Gamma(p)},$$

we obtain

$$\begin{aligned} \zeta_2(s) &= \zeta_R(2s-1) \frac{\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s)} \int_a^b f(x)^{2s-1} \sqrt{1+f'(x)^2} dx \\ &\quad + \zeta_R(2s+1) \frac{\Gamma(s+\frac{1}{2})}{\sqrt{\pi}\Gamma(s)} s \left[\frac{5s+1}{3} \int_a^b \frac{f(x)^{2s-1} f'(x)^2}{\sqrt{1+f'(x)^2}} dx + \int_a^b \frac{f(x)^{2s} f''(x)}{(1+f'(x)^2)^{3/2}} dx \right] + \dots \\ &\quad - \frac{1}{2} \zeta_R(2s) [f(a)^{2s} + f(b)^{2s}] \\ &\quad - \zeta_R(2s+2) f(a)^{2s} \frac{s^2}{4} \left[\frac{f'^2(a)}{1+f'^2(a)} \frac{5s+3}{4} + \frac{f''(a)f(a)}{(1+f'^2(a))^2} \right] \\ &\quad - \zeta_R(2s+2) f(b)^{2s} \frac{s^2}{4} \left[\frac{f'^2(b)}{1+f'^2(b)} \frac{5s+3}{4} + \frac{f''(b)f(b)}{(1+f'^2(b))^2} \right] - \dots \end{aligned}$$

For $s = 1, \frac{1}{2}, 0, -\frac{1}{2}$, etc, the residue or value of $\zeta_2(s)$ may have two contributions from the series above, since both $\zeta_R(s)$ and $\Gamma(s)$ have poles. For example, using properties of the

Riemann zeta function, $\zeta_R(0) = -\frac{1}{2}$ and $\text{Res } \zeta_R(1) = 1$, we have

$$\text{Res}\zeta_2(1) = \frac{1}{2} \int_a^b f(x) \sqrt{1 + f'(x)^2} dx, \quad (35)$$

$$\text{Res}\zeta_2\left(\frac{1}{2}\right) = -\frac{1}{2\pi} \int_a^b \sqrt{1 + f'(x)^2} dx - \frac{f(a) + f(b)}{4}, \quad (36)$$

$$\zeta_2(0) = \frac{1}{2}, \quad (37)$$

$$\begin{aligned} \text{Res}\zeta_2\left(-\frac{1}{2}\right) = & -\frac{1}{2\pi} \int_a^b \left[-\frac{1}{8} \frac{f'^2}{f^2 \sqrt{1 + f'^2}} + \frac{1}{4} \frac{f''}{f(1 + f'^2)^{3/2}} \right] dx \\ & - \frac{1}{32f(a)} \left[\frac{1}{8} \frac{f'^2(a)}{1 + f'^2(a)} + \frac{f''(a)f(a)}{(1 + f'^2(a))^2} \right] - \frac{1}{32f(b)} \left[\frac{1}{8} \frac{f'^2(b)}{1 + f'^2(b)} + \frac{f''(b)f(b)}{(1 + f'^2(b))^2} \right]. \end{aligned} \quad (38)$$

Adding up $\zeta_1(s)$ and $\zeta_2(s)$, and recalling that $\zeta_1(1)$ is finite, we obtain the following theorem for the full zeta function.

Theorem 4.1. *For the full zeta function of the Laplacian on a surface of revolution,*

$$\text{Res}\zeta(1) = \frac{1}{2} \int_a^b f(x) \sqrt{1 + f'(x)^2} dx. \quad (39)$$

$$\text{Res}\zeta\left(\frac{1}{2}\right) = -\frac{f(a) + f(b)}{4}. \quad (40)$$

$$\zeta(0) = 0. \quad (41)$$

$$\text{Res}\zeta\left(-\frac{1}{2}\right) = -\frac{1}{32f(a)} \left[\frac{1}{8} \frac{f'^2(a)}{1 + f'^2(a)} + \frac{f''(a)f(a)}{(1 + f'^2(a))^2} \right] - \frac{1}{32f(b)} \left[\frac{1}{8} \frac{f'^2(b)}{1 + f'^2(b)} + \frac{f''(b)f(b)}{(1 + f'^2(b))^2} \right]. \quad (42)$$

The geometric interpretation of the formulas above will be discussed in Section 5.

Derivative of $\zeta_2(s)$ at $s = 0$

The derivative of $\zeta_2(s)$ at 0 can be calculated in a similar but more subtle way. The following theorem is the main result of the paper.

Theorem 4.2. *The derivative at zero of the holomorphic extension of the zeta function on the two-dimensional surface of revolution is given by*

$$\zeta'(0) = -2 \ln \phi \left(e^{-2\frac{\pi^2}{A}} \right) + \frac{\pi^2}{6A} + \frac{1}{6} \int_a^b \frac{f'(x)^2}{f(x) \sqrt{1 + f'(x)^2}} dx + \frac{1}{2} \int_a^b \frac{f''(x)}{(1 + f'(x)^2)^{3/2}} dx, \quad (43)$$

where A is defined in Eq. (27), and ϕ is the Euler function [2, 37] defined by

$$\phi(q) = \prod_{k=1}^{\infty} (1 - q^k). \quad (44)$$

Proof. Integrating Eq. (29) by parts, we find that

$$\zeta_2(s) = 2s \frac{\sin \pi s}{\pi} \sum_{k=1}^{\infty} k^{-2s} \int_0^{\infty} z^{-s-1} \ln D_k(-k^2 z) dz. \quad (45)$$

$S_1(k, 0; x)$ and $\phi_k(0; b)$ can be computed exactly,

$$S_1(k, 0; x) = \frac{\sqrt{1 + f'(x)^2}}{f(x)} k, \quad \phi_k(0; b) = \frac{f(a)}{\sqrt{1 + f'(a)^2}} \frac{e^{kA} - e^{-kA}}{2k},$$

where A is defined by Eq. (27). This allows us to write

$$\ln D_k(-k^2 z) = \int_a^b S_1(k, z; x) dx - Ak - \frac{1}{2} \ln \frac{S_1(k, z; a) S_1(k, z; b)}{S_1(k, 0; a) S_1(k, 0; b)} + \ln \frac{1 - e^{-2 \int_a^b S_1(k, z; x) dx}}{1 - e^{-2Ak}}. \quad (46)$$

By the asymptotic expansion in Eq. (30), we write $S_1(k, z; x)$ as

$$S_1(k, z; x) = s_{-1}(z; x)k + \frac{s_1(z; x)}{k} + R(k, z; x),$$

where the remainder $R(k, z; x) = O(y^{-3/2}) = O(k^{-3}z^{-3/2})$ as $y \rightarrow \infty$. In addition, since $S_1(k, 0; x) = s_{-1}(0; x)k$, we have $s_1(z; x) = O(z)$ and $R(k, z; x) = O(k^{-3}z)$ as $z \rightarrow 0$. It implies that $\ln S_1(k, z; x)$ can be written as

$$\ln S_1(k, z; x) = \ln(s_{-1}(z; x)k) + D(k, z; x),$$

where $D(k, z; x) = O(y^{-1}) = O(k^{-2}z^{-1})$ as $y \rightarrow \infty$, and $D(k, z; x) = O(k^{-2}z)$ as $z \rightarrow 0$. Substituting Eq. (46) into Eq. (45) and using the expressions for $S_1(k, z; x)$, we have

$$\begin{aligned} \zeta_2(s) = 2s \frac{\sin \pi s}{\pi} \sum_{k=1}^{\infty} k^{-2s} \int_0^{\infty} z^{-s-1} \left\{ \int_a^b \left[\frac{\sqrt{1 + f'^2}}{f} k(\sqrt{t+1} - 1) + \frac{s_1(z; x)}{k} + R(k, z; x) \right] dx \right. \\ \left. - \frac{\ln(zf(a)^2 + 1) + \ln(zf(b)^2 + 1)}{4} - \frac{D(k, z; a) + D(k, z; b)}{2} + \ln \frac{1 - e^{-2 \int_a^b S_1(k, z; x) dx}}{1 - e^{-2Ak}} \right\} dz. \end{aligned}$$

The asymptotics of $R(k, z; x)$ and $D(k, z; x)$ as $y \rightarrow \infty$ and as $z \rightarrow 0$ implies that

$$\sum_{k=1}^{\infty} k^{-2s} \int_0^{\infty} z^{-s-1} dz \left\{ \int_a^b R(k, z; x) dx - \frac{D(k, z; a) + D(k, z; b)}{2} \right\}$$

is analytic for $-\frac{1}{2} < \Re s < 1$. Therefore,

$$\begin{aligned} \zeta_2(s) = 2s \frac{\sin \pi s}{\pi} \sum_{k=1}^{\infty} k^{-2s} \int_0^{\infty} z^{-s-1} \left\{ \int_a^b \left[\frac{\sqrt{1 + f'^2}}{f} k(\sqrt{t+1} - 1) + \frac{s_1(z; x)}{k} \right] dx \right. \\ \left. - \frac{\ln(zf(a)^2 + 1) + \ln(zf(b)^2 + 1)}{4} + \ln \frac{1 - e^{-2 \int_a^b S_1(k, z; x) dx}}{1 - e^{-2Ak}} \right\} dz + O(s^2). \end{aligned}$$

We now calculate each one of these terms individually. For the first one, we have

$$\begin{aligned}
& 2s \frac{\sin \pi s}{\pi} \sum_{k=1}^{\infty} k^{-2s} \int_0^{\infty} z^{-s-1} dz \int_a^b \frac{\sqrt{1+f'(x)^2}}{f(x)} k(\sqrt{t+1}-1) dx \\
&= 2 \frac{\sin \pi s}{\pi} \sum_{k=1}^{\infty} k^{1-2s} \int_a^b f(x)^{2s} \frac{\sqrt{1+f'(x)^2}}{f(x)} dx \int_0^{\infty} \frac{t^{-s}}{2\sqrt{t+1}} dt \\
&= \zeta_R(2s-1) \frac{\Gamma(s-\frac{1}{2})}{\Gamma(\frac{1}{2})\Gamma(s)} \int_a^b f(x)^{2s-1} \sqrt{1+f'(x)^2} dx \\
&= s \frac{A}{6} + O(s^2),
\end{aligned}$$

where we used $\zeta_R(-1) = -1/12$.

For the second term, substituting $s_1(z; x)$ from Eq. (31),

$$\begin{aligned}
& 2s \frac{\sin \pi s}{\pi} \sum_{k=1}^{\infty} k^{-2s} \int_0^{\infty} z^{-s-1} dz \int_a^b \frac{s_1(z; x)}{k} dx \\
&= 2s \frac{\sin \pi s}{\pi} \sum_{k=1}^{\infty} k^{-2s-1} \int_a^b \frac{f^{2s}(x) dx}{\sqrt{1+f'(x)^2}} \int_0^{\infty} \frac{t^{-s-1} dt}{\sqrt{t+1}} \left[-\frac{t(t-4)}{8(t+1)^2} \frac{f'(x)^2}{f(x)} + \frac{1}{4} \frac{t}{t+1} \frac{f''(x)}{1+f'(x)^2} \right] \\
&= s \zeta_R(2s+1) \frac{\Gamma(s+\frac{1}{2})}{\Gamma(\frac{1}{2})\Gamma(s)} \left[\frac{5s+1}{3} \int_a^b \frac{f(x)^{2s} f'(x)^2}{f(x) \sqrt{1+f'(x)^2}} dx + \int_a^b \frac{f(x)^{2s} f''(x)}{(1+f'(x)^2)^{3/2}} dx \right] \\
&= s \left[\frac{1}{6} \int_a^b \frac{f'(x)^2}{f(x) \sqrt{1+f'(x)^2}} dx + \frac{1}{2} \int_a^b \frac{f''(x)}{(1+f'(x)^2)^{3/2}} dx \right] + O(s^2).
\end{aligned}$$

For the third term, we obtain

$$\begin{aligned}
& 2s \frac{\sin \pi s}{\pi} \sum_{k=1}^{\infty} k^{-2s} \int_0^{\infty} z^{-s-1} \left[-\frac{\ln(zf(a)^2+1) + \ln(zf(b)^2+1)}{4} \right] dz \\
&= -\frac{1}{2} \frac{\sin \pi s}{\pi} \sum_{k=1}^{\infty} k^{-2s} \int_0^{\infty} \frac{t^{-s} dt}{t+1} [f(a)^{2s} + f(b)^{2s}] \\
&= -\frac{1}{2} \zeta_R(2s) [f(a)^{2s} + f(b)^{2s}] \\
&= \frac{1}{2} + s \left[-2\zeta'_R(0) + \frac{\ln f(a) + \ln f(b)}{2} \right] + O(s^2) \\
&= \frac{1}{2} + s \left[\ln(2\pi) + \frac{\ln f(a) + \ln f(b)}{2} \right] + O(s^2),
\end{aligned}$$

where we used $\zeta'_R(0) = -\frac{1}{2}\ln(2\pi)$. The fourth term involves the Euler function defined in Eq. (44),

$$\begin{aligned}
& 2s \frac{\sin \pi s}{\pi} \sum_{k=1}^{\infty} k^{-2s} \int_0^{\infty} z^{-s-1} \ln \frac{1 - e^{-2 \int_a^b S_1(k,z;x) dx}}{1 - e^{-2Ak}} dz \\
&= 2s \frac{\sin \pi s}{\pi} \left\{ \sum_{k=1}^{\infty} k^{-2s} \int_1^{\infty} z^{-s-1} \ln \frac{1 - e^{-2 \int_a^b S_1(k,z;x) dx}}{1 - e^{-2Ak}} dz + O(1) \right\} \\
&= -2s \frac{\sin \pi s}{\pi} \left\{ \sum_{k=1}^{\infty} k^{-2s} \int_1^{\infty} z^{-s-1} \ln(1 - e^{-2Ak}) dz + O(1) \right\} \\
&= -2s \sum_{k=1}^{\infty} \ln(1 - e^{-2Ak}) + O(s^2) \\
&= -2s \ln \phi(e^{-2A}) + O(s^2).
\end{aligned}$$

Adding these four terms gives

$$\begin{aligned}
\zeta'_2(0) &= -2 \ln \phi(e^{-2A}) + \frac{A}{6} + \ln(2\pi) + \frac{\ln f(a) + \ln f(b)}{2} \\
&\quad + \frac{1}{6} \int_a^b \frac{f'(x)^2}{f(x) \sqrt{1 + f'(x)^2}} dx + \frac{1}{2} \int_a^b \frac{f''(x)}{(1 + f'(x)^2)^{3/2}} dx.
\end{aligned} \tag{47}$$

Applying the following formula, which will be proved as Lemma 4.3,

$$\ln \phi(e^{-2\pi r}) - \ln \phi\left(e^{-\frac{2\pi}{r}}\right) = \frac{1}{2} \left[\frac{\pi}{6} \left(r - \frac{1}{r}\right) - \ln r \right],$$

with $r = A/\pi$, we have

$$\begin{aligned}
\zeta'_2(0) &= -2 \ln \phi(e^{-2\frac{\pi^2}{A}}) + \frac{\pi^2}{6A} + \ln(2A) + \frac{\ln f(a) + \ln f(b)}{2} \\
&\quad + \frac{1}{6} \int_a^b \frac{f'(x)^2}{f(x) \sqrt{1 + f'(x)^2}} dx + \frac{1}{2} \int_a^b \frac{f''(x)}{(1 + f'(x)^2)^{3/2}} dx.
\end{aligned} \tag{48}$$

We now add $\zeta'_1(0)$ from Eq. (26) with $\zeta'_2(0)$ above to obtain Eq. (43). \square

Lemma 4.3. *The following formula holds for the Euler function,*

$$\ln \phi(e^{-2\pi r}) - \ln \phi\left(e^{-\frac{2\pi}{r}}\right) = \frac{1}{2} \left[\frac{\pi}{6} \left(r - \frac{1}{r}\right) - \ln r \right]. \tag{49}$$

Proof. Consider the special case of $f \equiv R/\pi$ on an interval of length L . Then the eigenvalues are

$$\lambda_{k,n} = \left(\frac{k\pi}{R}\right)^2 + \left(\frac{n\pi}{L}\right)^2.$$

Due to the symmetry between R and L in the spectrum, $\zeta_2(s)(R, L) = \zeta_2(s)(L, R)$. Therefore, by Eq. (47),

$$\zeta'_2(0) = -2 \ln \phi\left(e^{-2\pi \frac{L}{R}}\right) + \frac{\pi}{6} \frac{L}{R} + \ln(2R) = -2 \ln \phi\left(e^{-2\pi \frac{R}{L}}\right) + \frac{\pi}{6} \frac{R}{L} + \ln 2L.$$

Letting $r = L/R$, we obtain Eq. (49). \square

5. DISCUSSION AND CONCLUSION

The geometric interpretation of the zeta function's residues and values can be found from the heat kernel expansion of the Laplacian on the surface of revolution; Refs. [27, 29, 43] gave extensive reviews of the heat kernel expansion. On a compact manifold, the heat kernel of the Laplacian, integrated over space, is defined as

$$\theta(t) = \sum_{\lambda_k \in \text{Spec}(\Delta)} e^{-t\lambda_k}.$$

For a two-dimensional manifold with boundary, denoted by Ω , the heat kernel has the following expansion,

$$\theta(t) = \frac{C_{-1}}{t} + \frac{C_{-\frac{1}{2}}}{\sqrt{t}} + C_0 + C_{\frac{1}{2}}\sqrt{t} + C_1 t + \dots$$

The coefficients are integrations of geometric invariants over Ω and its boundary $\partial\Omega$ [6]. For the surface of revolution, if the Dirichlet boundary condition is taken,

$$C_{-1} = \frac{|\Omega|}{4\pi}, \quad C_{-\frac{1}{2}} = -\frac{|\partial\Omega|}{8\sqrt{\pi}}, \quad C_0 = \frac{\chi}{6}, \quad C_{\frac{1}{2}} = \frac{1}{256\sqrt{\pi}} \int_{\partial\Omega} [k_g^2(\sigma) - 8k(\sigma)] d\sigma,$$

where $|\Omega|$ is the area of Ω , $|\partial\Omega|$ is the total length of $\partial\Omega$, χ is Euler characteristic of Ω , $k_g(\sigma)$ is the geodesic curvature of the boundary at σ , and $k(\sigma)$ is the Gaussian curvature of the surface at σ . For the surface of revolution, $\chi = 0$.

The zeta function is related to the heat kernel by

$$\zeta(s) = \frac{1}{\Gamma(s)} \int_0^\infty dt \theta(t) t^{s-1}.$$

Substituting the heat kernel expansion into the equation above, we have

$$\begin{aligned} \text{Res}\zeta(1) &= C_{-1} = \frac{|\Omega|}{4\pi} = \frac{1}{4\pi} \int_a^b 2\pi f(x) \sqrt{1 + f'(x)^2} dx, \\ \text{Res}\zeta\left(\frac{1}{2}\right) &= \frac{C_{-\frac{1}{2}}}{\sqrt{\pi}} = -\frac{|\partial\Omega|}{8\pi} = -\frac{2\pi f(a) + 2\pi f(b)}{8\pi}, \\ \zeta(0) &= C_0 = \frac{\chi}{6} = 0, \\ \text{Res}\zeta\left(-\frac{1}{2}\right) &= -\frac{C_{\frac{1}{2}}}{2\sqrt{\pi}} = -\frac{1}{512\pi} \int_{\partial\Omega} [k_g^2(\sigma) - 8k(\sigma)] d\sigma \\ &= -\frac{2\pi f(a)}{512\pi} \left[\frac{f'^2(a)}{f^2(a)(1 + f'^2(a))} + 8 \frac{f''(a)}{f(a)(1 + f'^2(a))^2} \right] \\ &\quad - \frac{2\pi f(b)}{512\pi} \left[\frac{f'^2(b)}{f^2(b)(1 + f'^2(b))} + 8 \frac{f''(b)}{f(b)(1 + f'^2(b))^2} \right]. \end{aligned}$$

They agree with Theorem 4.1 completely. It confirms the validity of the contour integral method extended to the surface of revolution. On the other hand, the determinant of the Laplacian, calculated by the contour integral method, cannot be derived from the heat kernel expansion directly.

Our results can also be benchmarked against an exactly solvable model with continuous spectrum. Consider the Laplacian on an infinitely long planar stripe of width L . Given the infinite volume of this configuration we need to introduce suitable zeta function densities.

To that end, we compactify the infinitely long planar stripe temporarily to a finite stripe of width $2\pi R$ and impose periodic boundary conditions along that direction. With Dirichlet boundary condition along the remaining direction, the spectrum is

$$\lambda_{k,n} = \left(\frac{k}{R}\right)^2 + \left(\frac{n\pi}{L}\right)^2, \quad k \in \mathbb{Z}, \quad n = 1, 2, 3, \dots$$

The associated zeta function is

$$\zeta(s) = \sum_{n=1}^{\infty} \sum_{k=-\infty}^{\infty} \left[\left(\frac{k}{R}\right)^2 + \left(\frac{n\pi}{L}\right)^2 \right]^{-s}.$$

As $R \rightarrow \infty$, which is the transition to the Riemann integral, the relevant zeta function density is

$$\begin{aligned} \zeta_c(s) &:= \lim_{R \rightarrow \infty} \frac{1}{R} \zeta(s) = \sum_{n=1}^{\infty} \int_{-\infty}^{\infty} \lambda_{nk}^{-s} dk \\ &= 2\zeta_R(2s-1) \int_0^{\infty} (x^2 + 1)^{-s} dx \left(\frac{L}{\pi}\right)^{2s-1} \\ &= \zeta_R(2s-1) B\left(\frac{1}{2}, s - \frac{1}{2}\right) \left(\frac{L}{\pi}\right)^{2s-1} \\ &= \sqrt{\pi} \zeta_R(2s-1) \frac{\Gamma(s - \frac{1}{2})}{\Gamma(s)} \left(\frac{L}{\pi}\right)^{2s-1}. \end{aligned}$$

In particular, we have

$$\text{Res} \zeta_c(1) = \frac{L}{2}, \quad \text{Res} \zeta_c\left(\frac{1}{2}\right) = -\frac{1}{2}, \quad \zeta_c(0) = 0, \quad \text{Res} \zeta_c\left(-\frac{1}{2}\right) = 0,$$

and

$$\zeta'_c(0) = \frac{\pi^2}{6L}.$$

The stripe can be regarded as a surface of revolution with $f(x) \equiv R$, and $R \rightarrow \infty$. For $f(x) \equiv R$, Theorem 4.1 gives

$$\text{Res} \zeta(1) = \frac{LR}{2}, \quad \text{Res} \zeta\left(\frac{1}{2}\right) = -\frac{R}{2}, \quad \zeta(0) = 0, \quad \text{Res} \zeta\left(-\frac{1}{2}\right) = 0,$$

and

$$\zeta'(0) = -2 \ln \phi \left(e^{-2\frac{\pi^2 R}{L}} \right) + \frac{\pi^2 R}{6L}.$$

One easily verifies, as the above computation shows, that these properties are recovered from

$$\zeta_c(s) = \lim_{R \rightarrow \infty} \frac{\zeta(s)}{R}.$$

As a conclusion, we have extended the contour integral method to the surface of revolution. By the WKB expansion, we calculated the zeta function at several important points. The results agree with those obtained from the heat kernel expansion. We also computed the determinant of the Laplacian on such a surface. The WKB expansion allows the evaluation of the zeta function at more points, however the expressions get increasingly more complicated. In the future, we can change the Dirichlet boundary condition to, say, Robin

boundary condition, and calculate the associated zeta function. We can also relax the condition that $f(x) > 0$ by allowing $f(a) = 0$ and/or $f(b) = 0$. This should allow the study of manifolds with cusps.

Acknowledgments

KK is supported by the National Science Foundation Grant PHY-0757791.

REFERENCES

- [1] L.V. Ahlfors. *Complex Analysis*. McGraw-Hill, Inc., New York, 1979.
- [2] T.M. Apostol. *Introduction to analytic number theory*. Springer Verlag, Berlin, 1976.
- [3] C. Bender. *Advanced mathematical methods for scientists and engineers I: Asymptotic methods and perturbation theory*. Springer, New York, 2010.
- [4] M. Bordag, G.L. Klimchitskaya, U. Mohideen, and V.M. Mostepanenko. *Advances in the Casimir effect*. Oxford Science Publications, 2009.
- [5] M. Bordag, U. Mohideen, and V.M. Mostepanenko. New developments in the Casimir effect. *Phys. Rept.*, 353:1–205, 2001.
- [6] T.P. Branson and P.B. Gilkey. The asymptotics of the Laplacian on a manifold with boundary. *Commun. Part. Diff. Equat.*, 15:245–272, 1990.
- [7] I.L. Buchbinder, S.D. Odintsov, and I.L. Shapiro. *Effective Action in Quantum Gravity*. Hilger, Bristol, 1992.
- [8] D. Burghelea, L. Friedlander, and T. Kappeler. On the determinant of elliptic differential and finite difference operators in vector bundles over S^1 . *Commun. Math. Phys.*, 138:1–18, 1991.
- [9] D. Burghelea, L. Friedlander, and T. Kappeler. Regularized determinants for pseudo differential in vector bundles over S^1 . *Integr. Equ. Oper. Theory*, 16:496–513, 1993.
- [10] D. Burghelea, T. Kappeler, P. McDonald, and L. Friedlander. On the functional logdet and related flows on the space of closed embedded curves on S^2 . *J. Funct. Anal.*, 120:440–466, 1994.
- [11] A.A. Bytsenko, G. Cognola, L. Vanzo, and S. Zerbini. Quantum fields and extended objects in space-times with constant curvature spatial section. *Phys. Rept.*, 266:1–126, 1996.
- [12] I. Chavel. *Eigenvalues in Riemannian Geometry*. Pure and Applied Mathematics, Vol. 115, Academic Press, New York, 1984.
- [13] G. Cognola, E. Elizalde, and K. Kirsten. Casimir energies for spherically symmetric cavities. *J. Phys. A: Math. Gen.*, 34:7311–7327, 2001.
- [14] S. Coleman. The fate of the false vacuum. 1. Semiclassical theory. *Phys. Rev.*, D15:2929–2936, 1977.
- [15] S. Coleman, V. Glaser, and A. Martin. Action minima among solutions to a class of Euclidean scalar field equations. *Commun. Math. Phys.*, 58:211–221, 1978.
- [16] J.S. Dowker and R. Critchley. Effective Lagrangian and energy momentum tensor in de Sitter space. *Phys. Rev.*, D13:3224–3232, 1976.
- [17] T. Dreyfuss and H. Dym. Product formulas for the eigenvalues of a class of boundary value problems. *Duke Math. J.*, 45:15–37, 1978.
- [18] G.V. Dunne and K. Kirsten. Functional determinants for radial operators. *J. Phys.*, A39:11915–11928, 2006.
- [19] G.V. Dunne and K. Kirsten. Simplified Vacuum Energy Expressions for Radial Backgrounds and Domain Walls. *J. Phys. A: Math. Theor.*, 42:075402, 2009.
- [20] E. Elizalde. *Ten Physical Applications of Spectral Zeta Functions*. Lecture Notes in Physics m35, Springer-Verlag, Berlin, 1995.
- [21] G. Esposito, A.Yu. Kamenshchik, and G. Pollifrone. *Euclidean Quantum Gravity on Manifolds with Boundary*. Fundamental Theories of Physics 85, Kluwer, Dordrecht, 1997.
- [22] R. Forman. Functional determinants and geometry. *Invent. Math.*, 88:447–493, 1987 Erratum, *ibid* 108 (1992) 453–454.
- [23] R. Forman. Determinants, finite-difference operators and boundary value problems. *Commun. Math. Phys.*, 147:485–526, 1992.

- [24] G. Fucci and K. Kirsten. Heat kernel coefficients for Laplace operators on the spherical suspension. *Commun. Math. Phys.*, to appear.
- [25] S. Gallot, D. Hulin, and J. Lafontaine. *Riemannian Geometry*. Springer, Berlin, 2004.
- [26] I.M. Gelfand and A.M. Yaglom. Integration in functional spaces and its applications in quantum physics. *J. Math. Phys.*, 1:48–69, 1960.
- [27] P.B. Gilkey. *Invariance Theory, the Heat Equation and the Atiyah-Singer Index Theorem*. CRC Press, Boca Raton, 1995.
- [28] S.W. Hawking. Zeta function regularization of path integrals in curved space-time. *Commun. Math. Phys.*, 55:133–148, 1977.
- [29] K. Kirsten. *Spectral Functions in Mathematics and Physics*. Chapman&Hall/CRC, Boca Raton, FL, 2002.
- [30] K. Kirsten and P. Loya. Computation of determinants using contour integrals. *Am. J. Phys.*, 76:60–64, 2008.
- [31] K. Kirsten and A.J. McKane. Functional determinants by contour integration methods. *Ann. Phys.*, 308:502–527, 2003.
- [32] K. Kirsten and A.J. McKane. Functional determinants for general Sturm-Liouville problems. *J. Phys. A: Math. Gen.*, 37:4649–4670, 2004.
- [33] N. Lebedev. *Special functions and their applications*. Dover Publications, Mineola, New York, 1972.
- [34] S. Levit and U. Smilansky. A theorem on infinite products of eigenvalues of Sturm-Liouville type operators. *Proc. Am. Math. Soc.*, 65:299–302, 1977.
- [35] P.D. Miller. *Applied asymptotic analysis*. American Mathematical Society, Providence, Rhode Island, 2006.
- [36] K.A. Milton. *The Casimir Effect: Physical Manifestations of Zero-Point Energy*. River Edge, USA: World Scientific, 2001.
- [37] NIST Digital Library of Mathematical Functions. dlmf.nist.gov. 27.14.2.
- [38] K. Okikiolu and C. Wang. Hessian of the zeta function of the Laplacian on forms. *Forum Math.*, 17:105–131, 2005.
- [39] B. Osgood, R. Phillips, and P. Sarnak. Extremals of determinants of Laplacians. *J. Funct. Anal.*, 80:148–211, 1988.
- [40] D.B. Ray and I.M. Singer. R-torsion and the Laplacian on Riemannian manifolds. *Advances in Math.*, 7:145–210, 1971.
- [41] R. Schoen and S.T. Yau. *Lectures on Differential Geometry*. International Press, Boston, 1994.
- [42] L.S. Schulman. *Techniques and Applications of Path Integration*. Wiley-Interscience, New York, 1981.
- [43] D.V. Vassilevich. Heat kernel expansion: User’s manual. *Phys. Rept.*, 388:279–360, 2003.

DEPARTMENT OF MATHEMATICS, STATISTICS, AND PHYSICS, WICHITA STATE UNIVERSITY, 1845 FAIRMOUNT, BOX 33, WICHITA, KS 67260-0033, USA

E-mail address: jeffres@math.wichita.edu

DEPARTMENT OF MATHEMATICS, BAYLOR UNIVERSITY, WACO, TX 76798, USA

E-mail address: Klaus.Kirsten@baylor.edu

DEPARTMENT OF MATHEMATICS, STATISTICS, AND PHYSICS, WICHITA STATE UNIVERSITY, 1845 FAIRMOUNT, BOX 33, WICHITA KS 67260-0033, USA

E-mail address: lu@math.wichita.edu